

Crystal Data: Orthorhombic. *Point Group:* $2/m\ 2/m\ 2/m$. Crystals are long prismatic, to 6 cm; terminated crystals are uncommon.

Physical Properties: *Cleavage:* Two directions, observed microscopically, may be parting. *Tenacity:* Brittle. *Fracture:* Uneven. Hardness = 6.5 VHN = 731-845 D(meas.) = 2.968 D(calc.) = 3.194

Optical Properties: Opaque, translucent on thin edges. *Color:* Dark brown to blue-green. *Luster:* Vitreous, greasy on fracture surfaces. *Optical Class:* Biaxial (-). $\alpha = 1.707$ $\beta = 1.745$ $\gamma = 1.776$ $2V(\text{meas.}) = 82^\circ$ *Pleochroism:* Distinct; X = pale yellow; Y = yellow; Z = brownish yellow. *Absorption:* $Z > Y \geq X$. *Dispersion:* $r < v$, strong.

Cell Data: Space Group: *Imma*. $a = 8.1538(4)$ $b = 10.5569(5)$ $c = 13.9882(6)$ $Z = 4$

X-ray Powder Pattern: Wolgidee Hills, Western Australia; nearly identical to batisite. 2.911 (100), 3.201 (80), 3.399 (75), 2.634 (70), 2.191 (60), 2.102 (60), 1.680 (60)

Chemistry:	(1)	(2)		(1)	(2)
SiO ₂	40.61	40.57	MnO	0.04	
TiO ₂	17.91	18.87	(Nb,Ta) ₂ O ₅	10.44	10.49
ZrO ₂	1.23		MgO	0.26	
Na ₂ O	5.82	5.59	CaO	0.82	0.20
K ₂ O	12.29	13.22	BaO	6.22	6.07
ZrO ₂		0.84	H ₂ O		0.73
Al ₂ O ₃	0.76		H ₂ O ⁺	0.54	
Fe ₂ O ₃	1.80	1.05	H ₂ O ⁻	0.10	
FeO	0.47		Total	99.61	97.63

(1) Khibiny massif, Russia; corresponds to $(\text{K}_{1.23}\text{Na}_{1.10}\text{Ba}_{0.24}\text{Ca}_{0.08}\text{Mg}_{0.04})_{\Sigma=2.69}(\text{Ti}_{1.31}\text{Nb}_{0.44}\text{Fe}_{0.17}\text{Zr}_{0.06})_{\Sigma=1.98}\text{Si}_{3.96}[\text{O}_{13.6}(\text{OH})_{0.4}]_{\Sigma=14.00}$. (2) Mount Rasvumchorr, Khibiny massif, Russia; electron microprobe analysis, H₂O from crystal structure, Mn, Mg, Al, Ta, Sr, F, Cl not detected, corresponds to $(\text{K}_{0.66}\text{Ba}_{0.23}\text{Na}_{0.07}\text{Ca}_{0.02})_{\Sigma=0.98}\text{K}_{1.00}\text{Na}_{1.00}(\text{Ti}_{1.40}\text{Nb}_{0.47}\text{Fe}_{0.08}\text{Zr}_{0.04})_{\Sigma=1.99}(\text{O}_{1.52}\text{OH}_{0.48})_{\Sigma=2.00}[\text{Si}_4\text{O}_{12}]$.

Polymorphism & Series: Forms a series with batisite.

Occurrence: Very rare, in pegmatite in a differentiated alkalic massif as a hydrothermal pectolite-natrolite vein cutting leucite-normative kalsilite-nepheline rischorrite (Khibiny massif, Russia); in lamproite (Leucite Hills, Wyoming, USA).

Association: Natrolite, manganoan pectolite, albite, potassic feldspar, astrophyllite, strontian fluorapatite, titanite (Khibiny massif, Russia); priderite, jeppeite, wadeite, perovskite, phlogopite, richterite (Wolgidee Hills, Western Australia).

Distribution: From Mts. Rasvumchorr and Yukspor, Khibiny massif, Kola Peninsula, Russia. In the Wolgidee Hills, West Kimberley district, Western Australia. From Emmons Mesa, Zirkel Mesa, and Black Butte, Leucite Hills, Sweetwater Co., Wyoming, USA.

Name: For Academician Demitrii Ivanovich *Shcherbakov* (1893-1966), Russian mineralogist and geochemist, Institute of Geology of Ore Deposits, Petrology, Mineralogy, and Geochemistry, Moscow, Russia.

Type Material: A.E. Fersman Mineralogical Museum, Russian Academy of Sciences, Moscow, Russia; 57256.

References: (1) Es'kova, E.M. and M.E. Kazakova (1954) Shcherbakovite - a new mineral. Doklady Acad. Nauk SSSR, 99, 837-841 (in Russian). (2) (1955) Amer. Mineral., 40, 788 (abs. ref. 1). (3) Vlasov, K.A., Ed. (1966) Mineralogy of rare elements, v. II, 550-553. (4) Prider, R.T. (1965) Noonkanbahite, a potassic batisite from the lamproites of Western Australia. Mineral. Mag., 34, 403-404. Mineralogical Society of America Handbook of Mineralogy Revised 9/6/2013

405. (5) Mitchell, R.H. (1990) Shcherbakovite in leucite phlogopite lamproites from the Leucite Hills, Wyoming. *Mineral. Mag.*, 54, 645-646. (6) Uvarova, Y.A, E. Sokolava, F.C. Hawthorne, R.P. L'Hevedich, and R.H. Mitchell (2003) The crystal chemistry of shcherbakovite from the Khibina massif, Kola peninsula, Russia. *Can. Mineral.*, 41, 1193-1201. (7) (2004) *Amer. Mineral.*, 89, 1834 (abs. ref. 6). (8) Krivovichev, S.V., V.N. Yakovenchuk, and Ya.A. Pakhomovskiy (2004) Topology and symmetry of the titanosilicate framework in the crystal structure of shcherbakovite, $\text{Na}(\text{K},\text{Ba})_2(\text{Ti},\text{Nb})_2\text{O}_2[\text{Si}_4\text{O}_{12}]$. *Zapiski Vseross. Mineral. Obshch.* 133(3), 55-63 (in Russian, English abstract). (9) (2005) *Amer. Mineral.*, 90, 773 (abs. ref. 8).